Trinity College

Trinity College Digital Repository

Faculty Scholarship

2-2013

Second addendum to the synoptic review of red algal genera

Craig W. Schneider

Trinity College, craig.schneider.1@trincoll.edu

Michael J. Wynne

University of Michigan - Ann Arbor

Follow this and additional works at: <https://digitalrepository.trincoll.edu/facpub>

Part of the [Biology Commons](#)

Review

Craig W. Schneider* and Michael J. Wynne

Second addendum to the synoptic review of red algal genera

Abstract: A second addendum to Schneider and Wynne's (Schneider, C.W. and M.J. Wynne. 2007. A synoptic review of the classification of red algal genera a half a century after Kylin's "Die Gattungen der Rhodophyceen". Bot. Mar. 50: 197–249.) "Synoptic review" of red algal genera and their classification is presented, with an updating of names of new taxa at the generic level and higher. In the past few years, the hierarchy of some genera has changed due to new subfamilies, which are cited and referenced below. There have also been the descriptions of some higher taxa in the past few years.

Keywords: classification; genera; Kylin; red algae; Rhodoplantae.

*Corresponding author: Craig W. Schneider, Department of Biology, Trinity College, Hartford, CT 16106, USA, e-mail: cschneid@trincoll.edu
Michael J. Wynne: Department of Ecology and Evolutionary Biology and Herbarium, University of Michigan, Ann Arbor, MI 48109, USA

Introduction

Gene sequencing and the resulting construction of phylogenetic trees continue to add to our knowledge of red algal genera and the classification of the Rhodophyta. Such molecular-based phylogenies often have shown the need to recognize segregate entities, at times necessitating the description of new genera or the resurrection of older generic names that had come to be treated as taxonomic synonyms using criteria based on the classical morphology and reproduction. Following the publication of Schneider and Wynne's (2007) "Synoptic review...of red algal genera," and our first addendum to it (Wynne and Schneider 2010), additional changes have appeared, leading us to issue this second addendum. Furthermore, we continue to find obscure valid names that we had overlooked in the synoptic review. We will periodically provide further addenda as sufficient new published information on red algal genera and classification appears.

The flood of new data coming in from the gene-sequence analyses is not only revealing the cases where some genera need to be split up and where others were cases of cryptic speciation but also where some genera and even families have been placed in orders where they clearly no longer belong. The red algal tree of life depicted by Verbruggen et al. (2010, figure 2), reconstructed using the Bayesian phylogenetic inference of DNA data mined from GenBank, reveals a contemporary view of our current understanding of the classification in the red algae at the class, subclass, order, and family levels. Their tree includes some taxa (*Atractophora*, Calosiphoniaceae, Inkyuleaceae, and Peyssonneliaceae), in which their traditional assignments are not supported by the present gene-sequence data.

With regard to the current realignment and descriptions of the many red algal genera due to molecular phylogenetics, it is worthwhile to refer to the ongoing discussion of the reliability of obtaining a DNA sequence data from the "archival" red algae. On one hand, some workers have questioned how successful the DNA extraction and sequencing can be for specimens older than about a dozen years (Saunders and McDevit 2012). On the other hand, other workers claim that the protocols have been successfully devised for the sequencing of the genes from the specimens going back into the early 19th century (Hughey and Gabrielson 2012). Confirmation of the results from the archival specimens with the DNA obtained from the recent collections from the same type localities should assist in resolving these questions.

Format of the list

The same format that was employed in the previous synoptic review and the first addendum (Schneider and Wynne 2007, Wynne and Schneider 2010) is followed in this second addendum. All the genera established after Kylin (1956) and added here have the authorities who described them, as well as the year and page of the protolog printed in bold face along with the proper names.

The “References” section contains the literature cited for the protologs of the genera since Wynne and Schneider (2010), as well as that not covered by Kylin (1956) and Schneider and Wynne (2007).

Kingdom Plantae

Phylum Rhodophyta

Subphylum Rhodellophytina

Class Rhodellophyceae

Order Dixonellales

Family Dixonellaceae

Note: Wynne and Schneider (2010) considered this family a synonym of the Glaucosphaeraceae based upon the rule of nomenclatural priority. Since then, however, the Glaucosphaeraceae was emended by Scott et al. (2011, 2012) to include only *Glaucosphaera* and transferred to the new order Glaucosphaerales [below]. This realignment of the Glaucosphaeraceae excluded *Dixonella* and *Neorhodella* from the family, allowing the resurrection of the Dixonellaceae to accommodate them (Scott et al. 2011, 2012).

***Bulboplastis* Kushibiki, A. Yokoyama, M. Iwataki, J.A. West et Y. Hara 2012: 116.**

Note: *Bulboplastis* was considered a member of the Dixonellaceae when described by Kushibiki et al. (2012), and it would remain there given its phylogenetic association with *Dixonella* and *Neorhodella* despite *Glaucosphaera* being segregated into its own order (Scott et al. 2012).

Order Glaucosphaerales E.C. Yang, J.L. Scott, S.Y. Yoon et J.A. West 2012: 71.

Note: The initial proposal by Yang et al. in Scott et al. (2011) to establish this order was not valid due to the lack of the Latin diagnosis and designation of the type family. The requirement for a Latin diagnosis lapsed beginning January 1, 2012, and so Yang et al. in Scott et al. (2012), with a diagnosis in English, later validated the name of the order and designated the type family to be the Glaucosphaeraceae Skuja (1954).

Class Porphyridiophyceae

Order Porphyridiales

Family Porphyridiaceae

***Timpurckia* E.C. Yang, J.L. Scott et J.A. West 2010: 606.**

Subphylum Metarhodophytina

Class Compsopogonophyceae

Order Erythropeltidales

Family Erythrotrichiaceae

Erythrotrichia* Aresch. 1850: 435. *nom. cons.

Note: Zuccarello et al. (2011) showed that *Erythrotrichia sensu lato* comprised at least seven well-supported infra-generic lineages, and the authors recognized the difficulty of linking these genetic groupings with the more than two dozen species names currently associated with the genus. They observed that the morphology and, at times, overlapping geographic locations of their isolates were not useful in tying these cryptic species to the historical binomials. Two species of *Erythrotrichia* were transferred to *Porphyrostromium*.

Subphylum Eurhodophytina

Class Bangiophyceae

Order Bangiales

Family Bangiaceae

Porphyra* C. Agardh 1824: xxxii, 190. *nom. cons.

Note: Several genera have, over the years, been subsumed under the speciose *Porphyra* (Schneider and Wynne 2007, Wynne and Schneider 2010). With the advances of molecular genetics in taxonomy, the workers have begun to find distinctions of the species at the generic level, and two genera (see below) with a large number of taxa have been resurrected for the genetic groupings that segregate them from *Porphyra* (Sutherland et al. 2011).

***Boreophyllum* S.C. Lindstr., N. Kikuchi, M. Miyata et Neefus in Sutherland et al. 2011: 1140.**

***Clymene* W.A. Nelson in Sutherland et al. 2011: 1138.**

***Fuscolium* S.C. Lindstr. in Sutherland et al. 2011: 1139.**

***Lysithea* W.A. Nelson in Sutherland et al. 2011: 1139.**

***Miuraea* N. Kikuchi, S. Arai, G. Yoshida, J.A. Shin et M. Miyata in Sutherland et al. 2011: 1137.**

***Pyropia* J. Agardh 1899: 149.**

Note: Formerly considered a taxonomic synonym of *Porphyra* (Wynne and Schneider 2010), this genus was resurrected for more than 75 species, leaving it as the most speciose genus in the Bangiales (Sutherland et al. 2011).

***Wildemanina* G. De Toni 1890: 144, 148.**

Synonym: ***Diploderma* Kjellm. (1893) *nom. illeg.* (non Link 1816)**

Note: *Wildemanina* was resurrected by Sutherland et al. (2011) from *Porphyra* where it was considered a taxonomic synonym (Schneider and Wynne 2007).

Class Florideophyceae**Subclass Hildenbrandiophycidae****Order Nemaliales****Family Liagoraceae**

***Neoizziella* S.-M. Lin, S.-Y. Yuan et Huisman 2011: 253.**

***Macrocarpus* S.-M. Lin, S.-Y. Yuan et Huisman 2011: 258.**

Note: Lin et al. (2011) based *Macrocarpus* on *Liagora perennis* I.A. Abbott.

***Sinocladia* C.K. Tseng et W. Li 2005: 53 [Chinese], 163 [Latin].**

Note: Zeng (Tseng) et al. (2005) placed their new genus in the Dermonemataceae, a family considered by most workers as a synonym of the Liagoraceae (Huisman 2006).

***Yoshizakia* S.-M. Lin, Huisman et Payri 2013: 164.**

Order Palmariales**Family Rhodophysemataceae**

***Rhodophysema* Batters 1900: 377.**

Synonym: ***Halosacciocolax* S. Lund (1959) nom. inval.**

Note: Saunders and Clayden (2010) provided a valid epithet for the species known as *Halosacciocolax kjellmanii* S. Lund, an invalidly published entity, after finding it grouped with the genus *Rhodophysema* (Clayden and Saunders 2010) in their COI-5P phylogenetic analyses.

Subclass Corallinophycidae**Order Corallinales****Family Corallinaceae****Subfamily Corallinoideae**

***Corallina* L. 1758: 805.**

Synonyms: ***Marginosporum* (Yendo) Ganesan (1968); *Serraticardia* (Yendo) P.C. Silva (1957); *Titanephlum* Nardo (1834); *Yamadaia* [*Yamadaea*] Segawa (1955).**

Note: Hind and Saunders (2013) presented a molecular phylogenetic evidence to propose the merger of both *Marginosporum* and *Serraticardia* within *Corallina*. The spelling we used previously, “*Titanephlum*” (Schneider and Wynne 2007), should be *Titanephlum*. The genus *Yamadaia* was reduced to a synonym of *Corallina* by virtue of Martone et al. (2012) transferring the generitype species, *Y. melobesioides* Segawa, into *Corallina*. The only other species, *Y. americana* E.Y. Dawson et R.L. Steele, was transferred into *Chiharaea* (Martone et al. 2012). With regard to the spelling of the name, the original spelling was *Yamadaia*, and Johansen (1969) altered it to *Yamadaea*. We notice a conflict in the *International Code of Nomenclature for algae, fungi, and plants* (Melbourne Code) (<http://www.iapt-taxon.org/nomen/main.php>)

between Art. 60.1, which states that “The original spelling of a name or epithet is to be retained,” and Recommendation 60B.1(a), which states that for a generic name derived from a surname ending in a vowel, the letter “-a” is added, except for a name ending in “-a,” in which case “-ea” is added. We accede to the Article rather than the Recommendation and, thus, accept Segawa’s (1955) original spelling.

***Ellisolandia* K. Hind et G.W. Saunders 2013: 109.**

Note: This segregate genus was based on *Corallina elongata* J. Ellis et Solander (Hind and Saunders 2013).

***Johansenia* K. Hind et G.W. Saunders 2013: 111.**

Note: This segregate genus was based on *Serraticardia macmillanii* (Yendo) P.C. Silva following the transfer of the type species of *Serraticardia*, *S. maxima* (Yendo) P.C. Silva, to *Corallina*.

***Pachyarthron* Manza 1937: 45.**

Note: Sato et al. (2009) suggested that *Pachyarthron* should be subsumed under *Corallina* on comparisons of multiple gene sequences, including those of the generitype, *P. cretaceum* (Postels et Rupr.) Manza. We await a more formal presentation of these results. The genus *Pachyarthron* has a confused nomenclatural history with *Bossiella* (Schneider and Wynne 2007), and at present, the two genera are considered distinct (Wynne and Schneider 2010).

Subfamily Lithophylloideae

***Amphiroa* J.V. Lamour. 1812: 186.**

Synonym: ***Eurytion* (Decne.) Lindl. (1846)**

Note: This synonym was omitted in Schneider and Wynne (2007). The genus was first recognized as section *Eurytion* of *Amphiroa* by Decaisne (1842), before being elevated by Lindley (1846).

Subfamily Mastophoroideae

Note: Kato et al. (2011) revised this subfamily created by Setchell (1943) to include only the genera *Mastophora* and *Metamastophora*.

Subfamily Hydrolithoideae A. Kato et M. Baba 2011: 669.

Note: This new subfamily was created to include *Hydrolithon* (Kato et al. 2011).

Subfamily Porolithoideae A. Kato et M. Baba 2011: 669.

Note: At present, this new subfamily includes only *Porolithon* (Kato et al. 2011).

Subfamily Neogoniolithoideae A. Kato et M. Baba 2011: 669.

Note: This new subfamily was created to include *Neogoniolithon* (Kato et al. 2011).

Subclass Rhodymeniophycidae

Order Ceramiales

Family Delesseriaceae

***Hymenopsis* S.-M. Lin, W.A. Nelson et Hommersand 2012: 64.**

Note: This new genus is based on the new species *Hymenopsis heterophylla*, which had previously gone under the name of *Hymenena palmata* f. *marginata sensu* Kylin from New Zealand (Lin et al. 2012b).

***Paraglossum* (J. Agardh) J. Agardh 1898: 213.**

Note: Using morphological and molecular studies, Lin et al. (2012a) showed that *Paraglossum* should be reinstated to include certain Southern Hemisphere species that had been placed in *Delesseria*. Earlier, *Paraglossum* had been relegated as a junior synonym of *Delesseria* (Schneider and Wynne 2007). Lin et al. (2012) showed that *Paraglossum* and *Apoglossum* were closely related genera in the newly described tribe Apoglosseae of the Delesseriaceae.

Family Rhodomelaceae

***Erythrocytis* J. Agardh 1876: 638.**

Synonym: ***Ricardia* Derb. et Sol. 1856: 211.**

Note: The name *Ricardia* was inadvertently omitted by Kylin (1956) and Schneider and Wynne (2007). Silva (1952) has reviewed the nomenclatural history of the names *Erythrocytis* and *Ricardia*.

***Laurenciella* Cassano, Gil-Rodríguez, Senties, Díaz-Larrea, M.C. Oliveira et M.T. Fujii 2012: 354.**

Note: On the basis of molecular data alone, this genus was segregated from *Laurencia* by Cassano et al. (2012) to include *L. marilzae* Gil-Rodríguez, Senties, Díaz-Larrea, Cassano et M.T. Fujii.

***Lampisiphonia* H.-G. Choi, Díaz Tapia et Bárbara in Bárbara et al. 2013a: 138.**

***Symphyocolax* M.-S. Kim in Kim et Cho 2010: 106.**

Note: This genus is a parasite of *Symphyocladia* (Kim and Cho 2010).

Order Gelidiales

Family Gelidiaceae

***Gelidium* J.V. Lamour. 1813: 128. nom. cons.**

Synonym: ***Porphyroglossum* Kütz. (1847)**

Note: The merger of the monotypic genus *Porphyroglossum* into *Gelidium* was proposed by Kim et al. (2011a). Their initial binomial, *Gelidium zollingeri* (Kütz.) comb. nov., however, was a later homonym of that by Sonder (1854), and so they proposed the *nomen novum* *Gelidium indonesianum* K.M. Kim, G.S. Gerung et S.M. Boo (Kim et al. 2011b).

Order Gigartinales

Family Caulacanthaceae

***Sterrocladia* F. Schmitz 1893: 77.**

Note: A second species was added to this uniquely freshwater gigartinean genus. Using the comparative gene-sequence data, Sherwood et al. (2012) showed *Sterrocladia* to be in a basal position relative to the Gigartinaceae and Phyllophoraceae and may well represent a new family. However, they believed that fresh collections of the generic type species, *S. amnica* (Mont.) F. Schmitz, from northern South America, were needed for the final disposition.

Family Chondrymeniaceae Rodríguez-Prieto, G. Sartoni, S.-M. Lin et Hommersand 2013: in press.

Note: The genus *Chondrymenia* Zanardini had previously been included in the Sarcodiaceae of the order Plocamiales (Schneider and Wynne 2007). But on the basis of *rbcL* gene sequence analyses, Rodríguez-Prieto et al. (2013) demonstrated that the genus was more closely allied with the Cystocloniaceae-Solieriaceae complex and justified being separated as a distinct family in the order Gigartinales.

Family Cruoriaceae

***Pseudopolyides* Gallardo, Bárbara et Cremades in Bárbara et al. 2013b: 193.**

Family Gigartinaceae

***Psilophycus* W.A. Nelson, Leister et Hommersand 2011: 220.**

Note: Based upon the morphological and reproductive characteristics, Nelson et al. (2011) found this genus to be basal to the other genera in the family, containing unique features thus far not found in the other genera of the family.

Family Kallymeniaceae

Note: Based upon the multigene analyses, Clarkston and Saunders (2012) placed *Beringia*, *Erythrophyllum*,

Kallymeniopsis, and *Hommersandia* in the Kallymeniaceae rather than the Crossocarpaceae where they previously had been treated.

Ectophora 1876: 689.

Note: As pointed out by D'Archino et al. (2011), *Ectophora* has had a complicated taxonomic history with periods as a synonym of *Callophyllis* and others as a discrete generic entity. Using multiple genetic markers, D'Archino et al. (2011) demonstrated that it demanded status at the generic level.

***Salishia* Clarkston et G.W. Saunders 2012: 51.**

Note: Clarkston and Saunders (2012) based their new genus on *Pugetia firma* Kylin.

***Stauromenia* D'Archino, W.A. Nelson et Zuccarello 2012: 452.**

Family Phylloporaceae

***Arche stenogramma* C.W. Schneid., Chengsupanimit et G.W. Saunders 2011: 447.**

Note: Along with a new species described from Bermuda, Schneider et al. (2011) included *Leptofaucha brasiliense* A.B. Joly in their new genus, a species that Dalen and Saunders (2007) had excluded from the Faucheaceae (Rhodymeniales).

***Coccotylus* Kütz. 1843**

Synonym: ***Ceratocolax*** Rosenv. (1898)

Note: Using the *rbcL* gene sequences and morphological investigations, Le Gall and Saunders (2010) found that the generitype and only species of the parasitic *Ceratocolax* was better assigned to *Coccotylus*, rendering the genus synonymous.

Family Schmitziellaceae Guiry, Garbary et G.W. Saunders in Guiry, 2012: 183

Note: Guiry et al. (in Guiry, 2012) described this new family, based on the genus *Schmitziella*, and provisionally assigned it to the order Gigartinales.

Order Peyssonneliales

Note: When pointing out the incorrect placement of the Peyssonneliaceae as a family of the Gigartinales, Verbruggen et al. (2010) were unaware of it already having been moved into its own order (Krayesky et al. 2009, Wynne and Schneider 2010).

Family Peyssonneliaceae

***Incendia* K.R. Dixon in Dixon and Saunders 2013: 84.**

Sonderophycus Denizot 1968: 206, 307.

Synonym: ***Sonderopelta*** Womersley et Sinkora (1981)

Note: Using both the chloroplast and nuclear DNA sequence data, Krayesky et al. (2009) demonstrated that the type species of *Sonderophycus* (*S. australis*) and *Sonderopelta* (*S. coriacea*) belonged in a single genus that was distinct from *Peyssonnelia*. However, they chose to recognize the junior name, *Sonderopelta*. Wynne (2011) pointed out that *Sonderophycus* has the priority over *Sonderopelta* and that the correct name for the type species is *Sonderophycus capensis* (Mont.) M.J. Wynne.

Order Halymeniales

Family Halymeniaceae

Dermocorynus P.L. Crouan et H.M. Crouan 1858: 69.

Note: Contrary to the findings of Wilkes et al. (2005), Gargiulo et al. (2013) resurrected *Dermocorynus* from the synonymy with *Grateloupia* using reproductive and molecular data.

***Glaphyrosiphon* Hommersand et Leister in Hommersand et al. 2010: 558.**

Grateloupia C. Agardh 1822: 221. *nom. cons.*

Synonym: ***Sinotubimorpha*** W.X. Li et Z.F. Ding (1998)

Note: Sheng et al. (2012) critically assessed the morphological features of two closely related red algal genera, *Grateloupia* and *Sinotubimorpha*, and found that it was difficult to define an obvious distinction between the two genera based only on the morphological differences. Using the *rbcL* sequence analysis, *Sinotubimorpha* formed a single monophyletic subclade within the large *Grateloupia* clade of the Halymeniaceae (Sheng et al. 2012). It was, therefore, concluded that *Sinotubimorpha* should be retained in synonymy with *Grateloupia* where Wynne (2005) and Schneider and Wynne (2007) had placed it earlier based on the West Indian generitype of *Sinotubimorpha*, *Grateloupia porracea* Kütz.

Pachymeniopsis Y. Yamada ex S. Kawabata 1954: 67.

Note: After being subsumed into *Grateloupia* by Kawaguchi (1997), Gargiulo et al. (2013) resurrected this genus from synonymy using reproductive and molecular data.

Phyllymenia J. Agardh 1848: 47.

Note: In contrast to the merger of *Phyllymenia* with *Grateloupia* by De Clerck et al. (2005), Gargiulo et al. (2013)

resurrected this genus from synonymy using reproductive and molecular data.

Prionitis J. Agardh 1851: 185.

Note: Wang et al. (2001) proposed the merger of this genus into *Grateloupia*. Using reproductive and molecular data, Gargiulo et al. (2013) resurrected this genus from synonymy.

Spongophloea Huisman, De Clerck, Prud'homme et Borowitzka 2011: 12.

Order Plocamiales

Family Plocamiaceae

Plocamium J.V. Lamour. 1813: 137. *nom. cons.*

Note: Cremades et al. (2011) investigated the type of *Plocamium cartilagineum* (L.) P.S. Dixon as well as the other original material of the European species of the genus, and they came to different conclusions than those reported by Saunders and Lehmkuhl (2005). This allowed for a taxonomic revision of the generitype and a better understanding of the other early named species of *Plocamium*.

Order Sebdeniales

Family Sebdeniaceae

Lesleigha Kraft et G.W. Saunders 2011: 421.

Order Rhodymeniales

Family Fryeellaceae

Agardhinula De Toni 1897: 64.

Note: Based upon the tetrasporangial ontogeny and characteristics, Saunders and McDonald (2010) and Schneider et al. (2012) suggested that this genus might possibly be a member of the Fryeellaceae, rather than the Rhodymeniaceae, where it had previously been placed.

Family Rhodymeniaceae

Cresia Losada-Troche et D.L. Ballant 2010: 296.

Incertae sedis generalis

Lukinia Perestenko 1996: 129, 203.

Reingardia Perestenko 1996: 129, 203.

Note: Because these two genera were first described directly after the treatment of the Phyllophoraceae (Gigartinales) in Perestenko (1996), we mistakenly attributed them to this family in our synoptic treatment of red algal genera (Schneider and Wynne 2007). It is unclear, at present, where they should be phylogenetically placed.

Acknowledgments: We thank Michael and Wendy Guiry for their continued work on AlgaeBase, an extremely valuable resource for the taxonomy and nomenclature of algae. We also thank Robert Andersen, Max Hommersand, Gary Saunders, John West and Gerry Kraft for assistance with this manuscript.

Received December 12, 2012; accepted January 22, 2013; online first February 27, 2013

References

- Bárbara, I., H.-G. Choi, A. Secilla, P. Díaz, J.M. Gorostiaga, T.-K. Seo, M.-Y. Jung and E. Berceibar. 2013a. *Lampisiphonia iberica* gen. et sp. nov. (Ceramiales, Rhodophyta) based on morphology and molecular evidence. *Phycologia* 52: 137–155.
- Bárbara, I., T. Gallardo, J. Cremades, R. Barreiro, I. Maneiro and G.W. Saunders. 2013b. *Pseudopolyides furcellarioides* gen. et sp. nov. (Gigartinales, Rhodophyta) an erect member of the Cruoriaceae based on morphological and molecular evidence. *Phycologia* 52: 191–203.
- Cassano, V., M.C. Oliveira, M.C. Gil-Rodríguez, A. Sentfies, J. Díaz-Larrea and M.T. Fujii. 2012. Molecular support for the establishment of the new genus *Laurenciella* within the *Laurencia* complex (Ceramiales, Rhodophyta). *Bot. Mar.* 55: 349–357.
- Clarkston, B.E. and G.W. Saunders. 2012. An examination of the red algal genus *Pugetia* (Kallymeniaceae, Gigartinales) with descriptions of *Salishia firma* gen. & comb. nov., *Pugetia cryptica* sp. nov. and *Beringia wynnei* sp. nov. *Phycologia* 51: 33–61.
- Clayden, S.L. and G.W. Saunders. 2010. Recognition of *Rubrointrusa membranacea* gen. et comb. nov., *Rhodonematella subimmersa* gen. et comb. nov. (with a reinterpretation of the life history) and the Meiodiscaceae fam. nov. within the Palmariales (Rhodophyta). *Phycologia* 49: 283–300.
- Cremades, J., R. Barreiro, I. Maneiro and G.W. Saunders. 2011. A new taxonomic interpretation of the type of *Plocamium cartilagineum* (Plocamiales, Florideophyceae) and its consequences. *Eur. J. Phycol.* 46: 125–142.
- Dalen, J.L. and G.W. Saunders. 2007. A review of the red algal genus *Leptofauchea* (Faucheaceae, Rhodymeniales) including a description of *L. chiloensis* sp. nov. *Phycologia* 46: 198–213.
- D'Archino, R., W.A. Nelson and G.C. Zuccarello. 2011. Diversity and complexity in New Zealand Kallymeniaceae (Rhodophyta): recognition of the genus *Ectophora* and description of *E. marginata* sp. nov. *Phycologia* 50: 241–255.
- D'Archino, R., W.A. Nelson and G.C. Zuccarello. 2012. *Stauromenia australis*, a new genus and species in the family Kallyme-

- niaceae (Rhodophyta) from southern New Zealand. *Phycologia* 51: 451–460.
- Decaisne, J. 1842. Essais sur une classification des algues et des polypiers calcifères de Lamouroux. Mémoire sur les corallines ou polypiers calcifère. *Ann. Sci. Nat., Bot., ser. 2*, 18: 96–128.
- De Clerck, O., B. Gavio, S. Fredericq, E. Cocquyt and E. Coppejans. 2005. Systematic reassessment of the red algal genus *Phyllymenia* (Halymeniaceae, Rhodophyta). *Eur. J. Phycol.* 40: 169–178.
- Dixon, K.R. and G.W. Saunders. 2013. DNA barcoding and phylogenetics of *Ramicrosta* and *Incendia* gen. nov., two early diverging lineages of the Peyssonneliaceae (Rhodophyta). *Phycologia* 52: 82–108.
- Gargiulo, G.M., M. Morabito and A. Manghisi. 2013. A re-assessment of reproductive anatomy and postfertilization development in the systematics of *Grateloupia* (Halymeniales, Rhodophyta). *Cryptogamie Algal.* 34: in press.
- Guiry, M.D. 2012. *A catalogue of Irish seaweeds*. A.R.G. Gantner Verlag, K.G., Ruggell, Liechtenstein. 250 pp.
- Hind, K.R. and G.W. Saunders. 2013. A molecular phylogenetic study of the tribe Corallineae (Corallinales, Rhodophyta) with an assessment of genus-level taxonomic features and descriptions of novel genera. *J. Phycol.* 49: 103–114.
- Hommersand, M.H., G.L. Leister, M.E. Ramírez, P.W. Gabrielson and W.A. Nelson. 2010. A morphological and phylogenetic study of *Glaphyrosiphon* gen. nov. (Halymeniaceae, Rhodophyta) based on *Grateloupia intestinalis* with descriptions of two new species: *Glaphyrosiphon lindaueri* from New Zealand and *Glaphyrosiphon chilensis* from Chile. *Phycologia* 49: 554–573.
- Hughey, J.R. and P.W. Gabrielson. 2012. Comment on “Acquiring DNA sequence data from dried archival red algae (Florideophyceae) for the purpose of applying available names to contemporary generic species: a critical assessment”. *Botany* 90: 1191–1194.
- Huisman, J.M. 2006. *Algae of Australia. Nemaliales*. Australian Biological Resources Study, Canberra and CSIRO Publishing, Melbourne. viii + 153 pp.
- Huisman, J.M., O. De Clerck, W.F. Prud’homme van Reine and M.A. Borowitzka. 2011. *Spongophloea*, a new genus of red algae based on *Thamnoclonium* sect. *Nematophorae* Weber-van Bosse (Halymeniales). *Eur. J. Phycol.* 46: 1–15.
- Johansen, H.W. 1969. Morphology and systematics of coralline algae with special reference to *Calliarthron*. *Univ. Calif. Publ. Bot.* 49: vi + 1–98.
- Kato, A., M. Baba and S. Suda. 2011. Revision of the Mastophoroideae (Corallinales, Rhodophyta) and polyphyly in nongeniculate species widely distributed on Pacific coral reefs. *J. Phycol.* 47: 662–672.
- Kawaguchi, S. 1997. Taxonomic notes on the Halymeniaceae (Gigartinales, Rhodophyta) from Japan. III. Synonymization of *Pachymeniopsis* Yamada in Kawabata with *Grateloupia* C. Agardh. *Phycol. Res.* 45: 9–21.
- Kim, K.M., G.S. Gerung and S.M. Boo. 2011a. Two-gene sequence and morphology of *Gelidium zollingeri* (Kützinger) comb. nov. (Gelidiales, Rhodophyta). *Algae* 26: 33–40.
- Kim, K.M., G.S. Gerung and S.M. Boo. 2011b. Corrigendum. Two-gene sequence and morphology of *Gelidium zollingeri* (Kützinger) comb. nov. (Gelidiales, Rhodophyta). *Algae* 26: 33–40, 209.
- Kim, M.S. and G.Y. Cho. 2010. A new red algal parasite *Symphycolax koreana* gen. et sp. nov. (Rhodomelaceae, Ceramiales), from Korea. *Algae* 25: 105–113.
- Kraft, G.T. and G.W. Saunders. 2011. Taxonomic and molecular studies of the family Sebdeniaceae (Sebdeniales, Rhodophyta): new species of *Lesleigha* gen. nov. and *Crassitegula* from Hawaii, east Asia and Lord Howe Island. *Eur. J. Phycol.* 46: 416–441.
- Kravesky, D.M., J.N. Norris, P.W. Gabrielson, D. Gabriel and S. Fredericq. 2009. A new order of red algae based on the Peyssonneliaceae, with an evaluation of the ordinal classification of the Florideophyceae (Rhodophyta). *Proc. Biol. Soc. Wash.* 122: 364–391.
- Kushibiki, A., A. Yokoyama, M. Iwataki, J. Yokoyama, J.A. West and Y. Hara. 2012. New unicellular red alga, *Bulboplastis apyrenoidosa* gen. et sp. nov. (Rhodellophyceae, Rhodophyta) from the mangroves of Japan: phylogenetic and ultrastructural observations. *Phycol. Res.* 60: 114–122.
- Kylin, H. 1956. *Die Gattungen der Rhodophyceen*. C.W.K. Gleerups, Lund. xv + 673 pp.
- Le Gall, L. and G.W. Saunders. 2010. DNA barcoding is a powerful tool to uncover algal diversity: a case study of the Phylloporaceae (Gigartinales, Rhodophyta) in the Canadian flora. *J. Phycol.* 46: 374–389.
- Lin, S.-M., S.-Y. Yang and J.M. Huisman. 2011. Systematics of *Liagora* with diffuse gonimoblasts based on *rbcl* sequences and carposporophyte development, including the description of the new genera *Neoizziella* and *Macrocarpus* (Liagoraceae, Rhodophyta). *Eur. J. Phycol.* 46: 249–262.
- Lin, S.-M., S. Fredericq and M.H. Hommersand. 2012a. Molecular phylogeny and developmental studies of *Apoglossum* and *Paraglossum* (Delesseriaceae, Rhodophyta) with a description of Apoglosseae trib. nov. *Eur. J. Phycol.* 47: 366–383.
- Lin, S.-M., W.A. Nelson and M.H. Hommersand. 2012b. *Hymenenopsis heterophylla* gen. et sp. nov. (Delesseriaceae, Rhodophyta) from New Zealand, based on a red alga previously known as *Hymenena palmata* f. *marginata* sensu Kylin, with emphasis on its cystocarp development. *Phycologia* 51: 62–73.
- Lin, S.-M., J.M. Huisman and C. Payri. 2013. Characterization of *Liagora ceranoides* (Liagoraceae, Rhodophyta) based on *rbcl* sequence analyses and carposporophyte development, including *Yoshizakia indopacifica* gen. et sp. nov. from the Indo-Pacific Region. *Phycologia* 52: 161–170.
- Lindley, J. 1846. *The vegetable kingdom; or, the structure, classification, and uses of plants, illustrated upon the natural system...with upwards of five hundred illustrations*. Published for the author by Bradbury and Evans, London. lxxviii + 908 pp.
- Lozada-Troche, C., D.L. Ballantine and H. Ruiz. 2010. *Cresia opalescens* gen. et sp. nov. (Rhodymeniaceae, Rhodophyta) from Puerto Rico, Caribbean Sea. *Cryptogamie Algal.* 31: 293–303.
- Martone, P.T., S.C. Lindstrom, K.A. Miller and P.W. Gabrielson. 2012. *Chiharaea* and *Yamadaia* (Corallinales, Rhodophyta) represent reduced and recently derived articulated coralline morphologies. *J. Phycol.* 48: 859–868.
- Nelson, W.A., G.L. Leister and M.H. Hommersand. 2011. *Psilophycus alveatus* gen. et comb. nov., a basal taxon in the Gigartinales (Rhodophyta) from New Zealand. *Phycologia* 50: 219–231.
- Perestenko, L.P. 1996 [1994]. *Red algae of the far-eastern seas of Russia*. Komarov Botanical Institute, Russian Academy of

- Sciences, St. Petersburg, and OLGA [a private publishing company]. 330 + [1] pp.
- Rodríguez-Prieto, C., G. Sartoni, S.-M. Lin and M.H. Hommersand. 2013. Comparative morphology and systematics of *Chondrymenia lobata* from the Mediterranean Sea and a phylogeny of the Chondrymeniaceae fam. nov. (Rhodophyta) based on *rbcl* sequence analyses. *Eur. J. Phycol.* 48: in press.
- Sato, D., S. Shimada, K. Kogame and M. Baba. 2009. A phylogenetic study of Japanese articulated coralline algae (Rhodophyta). *Phycologia* 48S: 115.
- Saunders, G.W. and K.V. Lehmkuhl. 2005. Molecular divergence and morphological diversity among four cryptic species of *Plocamium* (Plocamiales, Florideophyceae) in northern Europe. *Eur. J. Phycol.* 40: 293–312.
- Saunders, G.W. and B. McDonald. 2010. DNA barcoding reveals multiple overlooked Australian species of the red algal order Rhodymeniales (Florideophyceae), with resurrection of *Halopeltis* J. Agardh and description of *Pseudohalopeltis* gen. nov. *Botany* 88: 639–667.
- Saunders, G.W. and S.L. Clayden. 2010. Providing a valid epithet for the species widely known as *Halosaccicolax kjellmanii* S. Lund (Palmariales, Rhodophyta) – *Rhodophysema kjellmanii* sp. nov. *Phycologia* 49: 628.
- Saunders, G.W. and D.C. McDevit. 2012. Acquiring DNA sequence data from dried archival red algae (Florideophyceae) for the purpose of applying available names to contemporary genetic species: a critical assessment. *Botany* 90: 191–203.
- Schneider, C.W. and M.J. Wynne. 2007. A synoptic review of the classification of red algal genera a half a century after Kylin's "Die Gattungen der Rhodophyceen". *Bot. Mar.* 50: 197–249.
- Schneider, C.W., T. Chengsupanimit and G.W. Saunders. 2011. A new genus and species from the North Atlantic, *Archestenogramma profundum* from the North Atlantic, (Phylloporaceae, Rhodophyta), with taxonomic resolution of the orphaned *Leptofauchea brasiliensis*. *Eur. J. Phycol.* 46: 442–452.
- Schneider, C.W., D.W. Freshwater and G.W. Saunders. 2012. First report of *Halopeltis* (Rhodophyta, Rhodymeniaceae) from the non-tropical Northern Hemisphere: *H. adnata* (Okamura) comb. nov. from Korea, and *H. pellucida* sp. nov. and *H. willisii* sp. nov. from the North Atlantic. *Algae* 27: 95–108.
- Scott, J., E.C. Yang, J.A. West, A. Yokoyama, H.J. Kim, S.L. de Goër, C.J. O'Kelly, E. Orlova, S.Y. Kim, J.K. Park and H.S. Yoon. 2011. On the genus *Rhodella*, the emended orders Dixonellales and Rhodellales with a new order Glaucosphaerales (Rhodellophyceae, Rhodophyta). *Algae* 26: 277–288.
- Scott, J., E.C. Yang, J.A. West, A. Yokoyama, H.J. Kim, S.L. de Goër, C.J. O'Kelly, E. Orlova, S.Y. Kim, J.K. Park and H.S. Yoon. 2012. Corrigendum. On the genus *Rhodella*, the emended orders Dixonellales and Rhodellales with a new order Glaucosphaerales (Rhodellophyceae, Rhodophyta) (26: 277–288). *Algae* 27: 71–72.
- Segawa, S. 1955. Systematic anatomy of the articulated corallines. (Supplementary report). The structure and reproduction of *Yamadaia melobesioides* Segawa. *Bot. Mag. Tokyo* 68: 241–247.
- Setchell, W.A. 1943. *Mastophora* and the Mastophoreae: genus and subfamily of Corallinaceae. *Proc. Natl. Acad. Sci. USA* 29: 127–135.
- Sheng, Y.-W., W. Zhang, D. Zhao and H.-W. Wang. 2012. A morphological and molecular assessment of the genus *Sinotubimorpha* (Halymeniaceae, Rhodophyta). *J. Syst. Evol.* 50: 146–152.
- Sherwood, A.R., O. Necchi Jr., A.L. Carlile, H.D. Laughinghouse IV, S. Fredericq and R.G. Sheath. 2012. Characterization of a novel freshwater gigartinean red alga from Belize, with a description of *Sterrocladia belizeana* sp. nov. (Rhodophyta). *Phycologia* 51: 627–635.
- Silva, P.C. 1952. A review of nomenclatural conservation in the algae from the point of view of the type method. *Univ. Calif. Publ. Bot.* 25: 241–324.
- Sutherland, J.E., S.C. Lindstrom, W.A. Nelson, J. Brodie, M.D.J. Lynch, M.S. Hwang, H.-G. Choi, M. Miyata, N. Kikuchi, M.C. Oliveira, T. Farr, C. Neefus, A. Mols-Mortensen, D. Milstein, and K.M. Müller. 2011. A new look at an ancient order: generic revision of the Bangiales (Rhodophyta). *J. Phycol.* 47: 1131–1151.
- Verbruggen, H., C.A. Maggs, G.W. Saunders, L. Le Gall, H.S. Yoon and O. De Clerck. 2010. Data mining approach identifies research priorities and data requirements for resolving the red algal tree of life. *BMC Evol. Biol.* 10: 15 pp.
- Wang, H.W., S. Kawaguchi, T. Horiguchi and M. Masuda. 2001. A morphological and molecular assessment of the genus *Prionitis* J. Agardh (Halymeniaceae, Rhodophyta). *Phycol. Res.* 49: 251–261.
- Wilkes, R.J., L.M. McIvor and M.D. Guiry. 2005. Using *rbcl* sequence data to reassess the taxonomic position of some *Grateloupia* and *Dermocorynus* species (Halymeniaceae, Rhodophyta) from the north-eastern Atlantic. *Eur. J. Phycol.* 40: 53–60.
- Wynne, M.J. 2005. A checklist of benthic marine algae of the tropical and subtropical western Atlantic: second revision. *Nova Hedwigia, Beiheft* 129: ii + 1–152.
- Wynne, M.J. 2011. A checklist of benthic marine algae of the tropical and subtropical western Atlantic: third revision. *Nova Hedwigia, Beiheft* 140: [1]–166.
- Wynne, M.J. and C.W. Schneider. 2010. Addendum to the synoptic review of red algal genera. *Bot. Mar.* 53: 291–299.
- Yang, E.C., J. Scott, E. Orlova, D. Gauthier, F.C. Küpper, H.S. Yoon and U. Karsten. 2010. New taxa of Porphyridiophyceae (Rhodophyta): *Timspurckia oligopyrenoides* gen. et sp. nov. and *Erythrolobus madagascariensis* sp. nov. *Phycologia* 49: 604–616.
- Zeng, C. (Tseng, C.K.), R. Luan, Z. Chen and W. Li. 2005. *Flora algarum marinarum sinicarum Tomus II Rhodophyta No. II Acrochaetiales Nemaliales Bonnemaisoniales*. Science Press, Beijing. [i–vii], i–xxiii + 180 pp., pls I–VIII.
- Zuccarello, G.C., H.S. Yoon, H. Kim, L. Sun, S.L. de Goër and J.A. West. 2011. Molecular phylogeny of upright Erythropeltidales (Composopogonophyceae, Rhodophyta): multiple cryptic lineages of *Erythrotrichia carnea*. *J. Phycol.* 47: 627–637.